

# List of Tables

---

**Table 3.1** A summary of annual average particulate matter measurements at all air-quality monitoring stations for the Southern Appalachians.

**Table 3.2** A summary, by year, of the maximum 24-hour particulate matter measurements at all air-quality monitoring stations in the Southern Appalachians.

**Table 4.1** Percentage contribution by source category to pollutants which affect visibility in the eastern United States.

**Table 4.2** Median camera-based standard visual range (SVR) estimates, in miles (mi) and kilometers (km), and haziness values in deciview (dv), for the summer and winter seasons (for the combined years 1987-1993).

**Table 4.3** Annual camera-based standard visual range (SVR) estimates in miles (mi) and kilometers (km) and haziness values in deciview (dv), for the combined years 1987-1993.

**Table 4.4** Annual standard visual range (SVR) estimates in miles (mi) and kilometers (km) for combined years of sampling.

**Table 6.1** Listing of ozone exposure studies considered to identify the exposure effects levels for selected Southern Appalachian trees.

**Table 6.2** Ozone exposure levels associated with forest tree response.

**Table 6.3** Number of acres where ozone exposures may have been large enough to cause growth reductions.

**Table 6.4** Number of acres in each moisture index category.

**Table 6.5** Number of acres when ozone exposures and Palmer Hydrologic Index were favorable for growth loss.

**Table 6.6** Number of acres for each damage category when all years are combined.

